

Sequences and Sums

A sequence is a function t from a subset of the integers (usually N or Z^+) to a set S . In this discussion, the set S will be a set of numbers but we could have a sequence of colors, musical notes, or even computer programs.

We often write $a_n = t(n)$. Each a_n is a *term* of the sequence. The sequence a_1, a_2, a_3, \dots may be denoted by $\{a_n\}$ or $\{a_n\}_{n=1 \dots \infty}$. The numbers, $1, 2, 3, \dots$ are the term indices and a_1, a_2, a_3, \dots are the term values

1 Sequence Examples

Can you give the next term of each of these sequences? Can you give the function $t(n)$ that defines the sequence? Assume the first term is a_1 .

- 1, 1/2, 1/3, 1/4, 1/5, 1/6, ...
- 2, 4, 6, 8, 10, 12, ...
- 2, 4, 8, 16, 32, 64, ...
- 7, 8, 9, 10, 11, 12, ...
- 1/2, 1, 2, 4, 8, 16, ...
- 1, 3, 6, 10, 15, 21, ...
- 1, 1, 1, 1, 1, 1, ...
- 1, 1, 2, 3, 5, 8, ...
- 1/2, -2/3, 3/4, -4/5, 5/6, -6/7, ...
- 2, 3, 5, 7, 11, 13, ...

2 Some Special Sequences

2.1 Arithmetic Sequences

An *arithmetic sequence* is a sequence in which each term equals the preceding term plus a constant. A general arithmetic sequence looks like this $a, a + d, a + 2d, a + 3d, \dots$. The first term is a and the constant difference between terms is d .

What is the n^{th} term of the sequence $a, a + d, a + 2d, a + 3d, \dots$?

What is the next term and the n th term of each of these sequences?

- 4, 7, 10, 13, 16, 19, ...
- 7, -1, 5, 11, 17, 23, ...

2.2 Geometric Sequences

A *geometric sequence* is a sequence in which each term equals the preceding term times a constant. A general geometric sequence looks like this $a, ar, ar^2, ar^3, ar^4, \dots$. The first term is a and the constant ratio between successive terms is r .

What is the n^{th} term of the sequence $a, ar, ar^2, ar^3, ar^4, \dots$?

What is the next term and the n th term of each of these sequences?

2, 6, 18, 54, 162, ...
 1, -4, 16, -64, 256, ...

2.3 Quadratic Sequences

A *quadratic sequence* is a sequence whose n^{th} term is given by a quadratic function, $a_n = an^2 + b^n + c$. Here are some quadratic sequences. Can you find the quadratic function that generates them?

1, 4, 9, 16, 25, 36, ...
 6, 15, 28, 45, 66, 91, ...

An arithmetic sequence is given by a linear function and the difference between successive terms is a constant. In a quadratic sequence the differences between successive terms are given by a linear function and the second differences are constant. For example:

sequence	6	15	28	45	66	91	...
differences		9	13	17	21	25	...
second differences			4	4	4	4	...

3 Series and Partial Sums

A *series* is a sum of the terms of a sequence. Since a sequence has infinitely many terms, a series is the sum of infinitely many terms. We often sum only the first n terms of a sequence. The sum of the first n terms is called the n^{th} partial sum. The following is a sum of an infinite number of terms.

$$\sum_{k=1}^{\infty} a_k = a_1 + a_2 + a_3 + \dots$$

Below is the sum of the first n terms. This is the n^{th} *partial sum*. k is the *index of summation*; 1 is the *lower limit*; n is the *upper limit*.

$$\sum_{k=1}^n a_k = a_1 + a_2 + a_3 + \dots + a_n$$

This is also a partial sum. j is the index of summation; m is the lower limit; n is the upper limit.

$$\sum_{j=m}^n a_j = a_m + a_{m+1} + a_{m+2} + \dots + a_n$$

3.1 Arithmetic Sums

An *arithmetic sum* is a sum of terms of an arithmetic sequence.

$$\sum_{k=1}^n a + dk = (a + d) + (a + 2d) + (a + 3d) + \dots + (a + nd) = na + (1 + 2 + 3 \dots + n)d$$

$$na + \frac{n(n+1)}{2}d = n \left(\frac{a + 1 + a + n}{2} \right) = \text{num_terms} \left(\frac{\text{first_term} + \text{last_term}}{2} \right)$$

3.1.1 Examples

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2} \dots$$

$$1 + 2 + \dots + 100 = \frac{100(1+100)}{2} \dots = 5050$$

$$2 + 5 + 8 + 11 + 14 + 17 + 20 + 23 + 26 + 29 = \frac{10(2+29)}{2} \dots = 155$$

3.2 Geometric Sums and Series

A *geometric sum* is a sum of terms of a geometric sequence.

$$\sum_{k=1}^n ar^k = ar + ar^2 + ar^3 + \dots + ar^n$$

or, summing from 0

$$\sum_{k=0}^n ar^k = a + ar + ar^2 + ar^3 + \dots + ar^n$$

With geometric sequences, summing the infinite sequence may be meaningful. Infinite sums are called *series*.

$$\sum_{k=0}^{\infty} ar^k = a + ar + ar^2 + ar^3 + \dots$$

Let's find a formula for the n^{th} partial sum $\sum_{k=1}^n ar^k = ar + ar^2 + ar^3 + \dots + ar^n$.

$$\text{Let } S = \sum_{k=0}^n ar^k = a + ar + ar^2 + ar^3 + \dots + ar^n.$$

$$\text{Then } rS = \sum_{k=0}^n ar^{k+1} = ar + ar^2 + ar^3 + ar^4 + \dots + ar^n + ar^{n+1}.$$

$$S - rS = a - ar^{n+1} \text{ as all the other terms cancel out when we subtract } rS \text{ from } S.$$

$$\text{So } S = \frac{a - ar^{n+1}}{1 - r} = \frac{a(1 - r^{n+1})}{1 - r}.$$

3.2.1 Examples

$$\sum_{k=0}^6 \frac{1}{3^k} = 1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \frac{1}{3^4} + \frac{1}{3^5} + \frac{1}{3^6} = \frac{1 \left(1 - \frac{1}{3^7} \right)}{1 - \frac{1}{3}} = \frac{3^7 - 1}{3^7 - 3^6} = \frac{2187 - 1}{2197 - 729} = \frac{2186}{1468} \approx 1.499.$$

$$\sum_{k=0}^6 3^k = 1 + 3 + 3^2 + 3^3 + 3^4 + 3^5 + 3^6 = \frac{1(1 - 3^7)}{1 - 3} = \frac{2^7 - 1}{2} = \frac{2186}{2} = 1093.$$

$$\sum_{k=0}^6 \frac{1}{2^k} = 1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} + \frac{1}{2^5} + \frac{1}{2^6} = \frac{1 \left(1 - \frac{1}{2^7} \right)}{1 - \frac{1}{2}} = \frac{2^7 - 1}{2^7 - 2^6} = \frac{128 - 1}{128 - 64} = \frac{127}{64} \approx 1.984.$$

$$\sum_{k=0}^n \frac{1}{2^k} = \frac{1 \left(1 - \frac{1}{2^{n+1}} \right)}{1 - \frac{1}{2}} = \frac{1 - \frac{1}{2^{n+1}}}{\frac{1}{2}} = 2 - \frac{1}{2^n}. \text{ What happens when } n \text{ gets very large?}$$

$$\sum_{k=0}^{\infty} \frac{1}{2^k} = ?$$

3.3 Some Special Sums

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}.$$