

These notes are concerned with notational issues and relations. Throughout you will find “exercises.” You should work out solutions to the exercises that are not obvious to you, but you do not need to turn in solutions.

A warning. The notation I use differs from the notation you will see in the books we will use for the course. Being exposed to various notational conventions is a good thing, but may lead to confusion, so if something is not clear, please ask!

1 Initial Notation and Definitions

\mathbb{N} and ω both denote the natural numbers, *i.e.*, $\{0, 1, \dots\}$. The ordered pair whose first component is i and whose second component is j is denoted $\langle i, j \rangle$. $[i..j]$ denotes the closed interval $\{k \in \mathbb{N} : i \leq k \leq j\}$; parentheses are used to denote open and half-open intervals, *e.g.*, $[i..j)$ denotes the set $\{k \in \mathbb{N} : i \leq k < j\}$.

R is a *binary relation* on set S if $R \subseteq S \times S = \{\langle x, y \rangle : x, y \in S\}$. We abbreviate $\langle s, w \rangle \in R$ by sRw . A function is a relation such that xRy and xRw implies $y = w$.

Function application is sometimes denoted by an infix dot “.” and is left associative. That is, $f.x$ is the unique y such that xfy . This allows us to use the curried version of a function when it suits us, *e.g.*, we may write $f.x.y$ instead of $f(x, y)$. That is, $f.x.y$ is really $(f.x).y$, where f is a function of one argument that returns $f.x$, a function of one argument.

From highest to lowest binding power, we have: parentheses, function application, binary relations (*e.g.*, sBw), equality ($=$) and membership (\in), conjunction (\wedge) and disjunction (\vee), implication (\Rightarrow), and finally, binary equivalence (\equiv). Spacing is used to reinforce binding: more space indicates lower binding.

$\langle Qx : r : b \rangle$ denotes a quantified expression, where Q is the quantifier, x the bound variable, r the range of x (**true** if omitted), and b the body. We sometimes write $\langle Qx \in X : r : b \rangle$ as an abbreviation for $\langle Qx : x \in X \wedge r : b \rangle$, where r is **true** if omitted, as before.

Cardinality of a set S is denoted by $|S|$. $\mathcal{P}(S)$ denotes the powerset of S .

A function from $[0..n)$, where n is a natural number, is called a *finite sequence* or an *n-sequence*.

What are numbers as mathematical objects? von Neumann proposed the following: $0 = \emptyset$, $1 = \{0\}$, $2 = \{0, 1\}$, \dots , so $n = [0..n)$. Thus an n -sequence is a function from n .

An ω -sequence is a function from ω . We may sometimes refer to ω -sequences as infinite sequences, but as we will see there are infinite sequences that are “longer” than ω -sequences.

When we write $x \in \sigma$, for a sequence σ , we mean that x is in the range of σ .

2 Binary Relations

Let B, C be binary relations on set S . $B|_A$ denotes B *left-restricted* to the set A , i.e., $B|_A = \{\langle x, y \rangle : xBy \wedge x \in A\}$.

Some important definitions follow.

- B is *reflexive* if $\langle \forall x \in S :: xBx \rangle$.
- B is *irreflexive* if $\langle \forall x \in S :: \neg(xBx) \rangle$.
- B is *transitive* if $\langle \forall x, y, z \in S :: xBy \wedge yBz \Rightarrow xBz \rangle$.
- B is a *preorder* (also called a *quasi-order*) if it is reflexive and transitive.
- The identity relation, B^0 , is $\{\langle x, x \rangle : x \in S\}$.
- The *composition* of B and C is denoted $B;C$ and is the set $\{\langle b, c \rangle : \langle \exists x :: bBx \wedge xCc \rangle\}$.
- For all natural numbers i , B^{i+1} is $B^i;B$.

Exercise 1 Prove the following.

1. B is reflexive iff $B^0 \subseteq B$.
2. $B^1 = B$.
3. B is transitive iff $B^2 \subseteq B$.

We now continue with the definitions.

- B is *symmetric* if $\langle \forall x, y \in S :: xBy \Rightarrow yBx \rangle$.
- A preorder that is also symmetric is an *equivalence relation*.
- B is *asymmetric* if $\langle \forall x, y \in S :: xBy \Rightarrow \neg(yBx) \rangle$.
- B is *antisymmetric* if $\langle \forall x, y \in S :: xBy \wedge yBx \Rightarrow x = y \rangle$.
- A preorder that is antisymmetric is a *partial order*.
- If B is a partial order, $\langle S, B \rangle$ is a *poset*.
- The *inverse* of B is denoted B^{-1} and is $\{\langle x, y \rangle : yBx\}$.

Exercise 2 Prove the following.

1. B is symmetric iff $B^{-1} \subseteq B$.
2. B is antisymmetric iff $B \cap B^{-1} \subseteq B^0$.

If B is an equivalence relation, for each $x \in S$, it induces an *equivalence class* $[x]_B = \{y : xBy\}$. The *quotient* S/B is $\{[x]_B : x \in S\}$.

Exercise 3 Prove the following.

1. If B is an equivalence relation, then $[x]_B$ and $[y]_B$ are either identical or disjoint.
2. If C is a preorder, then
 - (a) $B = \{\langle x, y \rangle : xCy \wedge yCx\}$ is an equivalence relation.
 - (b) $\langle S/B, \preceq \rangle$ is a poset, where \preceq is defined as follows:

$$[x]_B \preceq [y]_B \equiv xCy.$$

We now continue with the definitions.

- B is *total* (also called *linear* or *connected*) if $\langle \forall x, y \in S :: xBy \vee yBx \rangle$.
- A *total order* is a partial order that is total.
- If B is a total order, $\langle S, B \rangle$ is a *toset*.
- An α -sequence $\langle a_0, a_1, a_2, \dots \rangle$, where $\alpha \in \omega \vee \alpha = \omega$, is *decreasing* in B if $\langle \forall i : i + 1 \in \alpha : a_{i+1}Ba_i \rangle$.
- B is *terminating* (also called *well-founded*) if there is no decreasing ω -sequence in B .
- If B is terminating, then $\langle S, B \rangle$ is a *well-founded structure*.
- The *strict part* of a relation B is $\{\langle x, y \rangle : xBy \wedge x \neq y\}$.
- B is a *strict partial order* if it is the strict part of some partial order. Strict total orders are defined in an analogous way.
- A *well order* is a strict total order that is well-founded.
- If B is a well order, $\langle S, B \rangle$ is a *woset*.
- For $T \subseteq S$:
 - If $(m \in T \wedge \langle \forall x \in T :: xBm \Rightarrow x = m \rangle)$, then m is a *minimal* element of T (under B).
 - If $(m \in T \wedge \langle \forall x \in T :: mBx \vee m = x \rangle)$, then m is the *least* element of T (under B).
 - If $(m \in S \wedge \langle \forall x \in T :: mBx \vee m = x \rangle)$, then m is a *lower bound* of T (under B).
 - The notions of *maximal*, *greatest*, and *upper bound* are defined dually, e.g., m is a maximal element of T under B iff m is a minimal element of T under B^{-1} .

Exercise 4 Prove the following.

1. B is total iff $B \cup B^{-1} = S \times S$.
2. B is a strict partial order iff it is irreflexive and transitive.
3. If \prec is a strict partial order and $x \preceq y \equiv x \prec y \vee x = y$ then \preceq is a partial order.
4. If \preceq is a preorder and $x \prec y \equiv x \preceq y \wedge \neg(y \prec x)$ then \prec is a strict partial order.
5. B is a strict total order iff
 - (a) B is irreflexive.
 - (b) B is transitive.
 - (c) $\langle \forall x, y \in S :: xBy \vee yBx \vee x = y \rangle$.
6. B is a well order iff it is well-founded and $\langle \forall x, y \in S :: xBy \vee yBx \vee x = y \rangle$.

Exercise 5 Let \prec be a strict partial order on S . Prove the following.

1. Prove that $\langle S, \prec \rangle$ is a well-founded structure iff all non-empty subsets of S have a minimal element under \prec .
2. Prove that $\langle S, \prec \rangle$ is a woset iff all non-empty subsets of S have a least element.

Given a set U (the “universe”), $X \subseteq U$, and a property P which is satisfied by some subsets of U , the P -sets, we say that C is the P -closure of X if C is the least P -set which includes X . If the P -sets include U and are closed under arbitrary intersections, we say that the P -sets of U form a *closure system*. If the P -sets of U form a closure system, then the P -closure of X always exists. It is $\bigcap \{Y \subseteq U : X \subseteq Y \wedge Y \text{ is a } P\text{-set}\}$.

Exercise 6 Prove the following, where $U = S \times S$.

1. The reflexive relations form a closure system.
2. The irreflexive relations do not form a closure system.
3. The symmetric relations form a closure system.
4. The asymmetric relations do not form a closure system.
5. The antisymmetric relations do not form a closure system.
6. The transitive relations form a closure system.

We can therefore speak of the reflexive closure, or the symmetric closure, or the transitive closure, or the reflexive, transitive closure, etc. B^+ denotes the transitive closure of B and B^* denotes the reflexive, transitive closure of B . This same notation is used in regular languages.